

SKEWADJOINT OPERATORS ON PSEUDOEUCCLIDEAN SPACES

Changrim Jang¹ and Phillip E. Parker

Mathematics Department
Wichita State University
Wichita KS 67260-0033
USA

crjang@math.twsu.edu

phil@math.twsu.edu

29 January 2003

Abstract: We give a complete classification in canonical forms on finite-dimensional vector spaces over $\mathbb{R}$.

MSC(1991): Primary 15A21; Secondary 15A57, 53C50.



¹On sabbatical leave; permanent address: School of Mathematics and Applied Physics, College of Natural Sciences, University of Ulsan, Ulsan 680-749, Republic of Korea. Email: crjang@mail.ulsan.ac.kr

1 Introduction

In this paper we give a complete classification of canonical forms for skewadjoint operators on pseudoeuclidean vector spaces, complementing O'Neill's results [7] on selfadjoint operators.

There is a certain skewadjoint operator in the integration of the geodesic equations on a 2-step nilpotent Lie group with a left-invariant metric tensor, usually called J after Eberlein [3]. In the Riemannian (positive definite) case, J^2 is always diagonalizable. This makes possible the completely explicit integration of the geodesic equations (compare [3] and [2]) and the rather complete and explicit determination of the conjugate loci (compare [5] and [6]). In attempting to extend some of these results to the pseudoriemannian (indefinite) case, it became apparent that we needed a set of canonical forms for such skewadjoint operators J . Obtaining the list turned out to be a sufficiently involved process that we are presenting it separately.

Throughout, we consider a (finite-dimensional) vector space V provided with an inner product $\langle, \rangle$; *i.e.*, a nondegenerate, symmetric bilinear form which is generally indefinite. When necessary, we comment on differences if the form is actually definite. Also, $J : V \rightarrow V$ will denote a skewadjoint linear operator.

We used [4] as our main reference for linear algebra, and [1] to a lesser extent, and refer to them for all details relevant to the computational procedures employed.

2 Preparation for the classification

To begin, we consider the minimal polynomial of J in the (preliminary) form

$$p(t) = \prod_{i=1}^p (t^2 - 2a_i t + (a_i^2 + b_i^2))^{k_i} \prod_{i=1}^q (t^2 + \lambda_i^2)^{l_i} \prod_{i=1}^r (t - \mu_i)^{m_i}$$

where $a_i \neq 0 \neq b_i$, $\lambda_i > 0$, and all $a_i, b_i, \lambda_i, \mu_i$ are real. The types of factors correspond to complex eigenvalues that are not pure imaginary, pure imaginary eigenvalues, and real eigenvalues, respectively.

Proposition 2.1 *For $\mu_i \neq 0$,*

$$\ker (J - \mu_i I)^{m_i} \quad \text{and} \quad \ker (J^2 - 2a_i J + (a_i^2 + b_i^2) I)^{k_i}$$

are null subspaces of V .

Proof: If $u, v \in \ker(J - \mu_i I)^{m_i}$, then

$$\begin{aligned}\langle u, (J + \mu_i I)^{m_i} v \rangle &= \langle (-J + \mu_i I)^{m_i} u, v \rangle \\ &= \langle (-1)^{m_i} (J - \mu_i I)^{m_i} u, v \rangle \\ &= \langle 0, v \rangle \\ &= 0.\end{aligned}$$

Note that $(J + \mu_i I)^{m_i} : \ker(J - \mu_i I)^{m_i} \rightarrow \ker(J - \mu_i I)^{m_i}$ is nonsingular. Thus it follows that $\langle u, v \rangle = 0$ for all u, v and $\ker(J - \mu_i I)^{m_i}$ is null as claimed. The other case is done similarly. $\square$

Proposition 2.2 *For nonzero μ_i , $\ker(J + \mu_i I)^{m_i}$ and $\ker(J^2 + 2a_i J + (a_i^2 + b_i^2)I)^{k_i}$ are complementary null subspaces to $\ker(J - \mu_i I)^{m_i}$ and to $\ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)$, respectively.*

Proof: Suppose $\ker(J + \mu_i I) = 0$. Then we see that $J + \mu_i I : W \rightarrow W$ is nonsingular, where

$$\begin{aligned}W &= \bigoplus_{i=1}^p \ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)^{k_i} \\ &\quad + \bigoplus_{i=1}^q \ker(J^2 + \lambda_i I)^{l_i} + \bigoplus_{j \neq i} \ker(J - \mu_j I)^{m_j}\end{aligned}$$

By the previous result, if $0 \neq u \in \ker(J - \mu_i I)$, then there exists $w \in W$ such that $\langle u, w \rangle = 1$. Since $J + \mu_i I$ is nonsingular on W , there exists $w' \in W$ such that $(J + \mu_i I)w' = w$. But now we have

$$\begin{aligned}1 &= \langle u, w \rangle = \langle u, (J + \mu_i I)w' \rangle \\ &= \langle (-J + \mu_i I)u, w' \rangle \\ &= -\langle (J - \mu_i I)u, w' \rangle \\ &= 0\end{aligned}$$

which is a contradiction. Thus a power term $(t + \mu_i)^{m'_i}$ must appear in $p(t)$. We shall show that $m'_i = m_i$.

Suppose $m'_i < m_i$. We can choose $u \in V$ such that $(J - \mu_i I)^{m_i-1}u \neq 0$ and $(J - \mu_i I)^{m_i}u = 0$. Then there exists $v \in W$ such that $\langle (J - \mu_i I)^{m_i-1}u, v \rangle = 1$. Since $m_i - 1 \geq m'_i$, $(-1)^{m_i-1}(J + \mu_i I)^{m_i-1}v \in W_1$ where

$$W_1 = \bigoplus_{i=1}^p \ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)^{k_i}$$

$$+ \bigoplus_{i=1}^q \ker (J^2 + \lambda_i I)^{l_i} + \bigoplus_{\mu_j \neq \pm \mu_i} \ker (J - \mu_j I)^{m_j}$$

Since $(J + \mu_i I)^{m_i} : W_1 \rightarrow W_1$ is nonsingular, there exists $v' \in W_1$ such that $(J + \mu_i)^{m_i} v' = (-1)^{m_i-1} (J + \mu_i I)^{m_i-1} v$. But now we have

$$\begin{aligned} 1 &= \langle u, (-1)^{m_i-1} (J + \mu_i I)^{m_i-1} v \rangle = \langle u, (J + \mu_i I)^{m_i} v' \rangle \\ &= \langle (-1)^{m_i} (J - \mu_i I)^{m_i} u, v' \rangle \\ &= 0 \end{aligned}$$

which is a contradiction. Thus $m'_i \geq m_i$. Interchanging roles, we get $m_i \geq m'_i$ so $m'_i = m_i$.

Now we show that $\ker(J + \mu_i I)^{m_i}$ is a complementary null subspace to $\ker(J - \mu_i I)^{m_i}$. For $0 \neq u \in \ker(J - \mu_i I)^{m_i}$, there exists $w \in W$ such that $\langle u, w \rangle = 1$. Now write $w = w_1 + w_2$ where $w_1 \in \ker(J + \mu_i I)^{m_i}$ and $w_2 \in W_1$. Since $(J + \mu_i I)^{m_i} : W_1 \rightarrow W_1$ is nonsingular, there exists w'_2 such that $(J + \mu_i I)^{m_i} w'_2 = w_2$. So we have

$$\begin{aligned} 1 &= \langle u, w \rangle = \langle u, w_1 + w_2 \rangle = \langle u, w_1 \rangle + \langle u, w_2 \rangle \\ &= \langle u, w_1 \rangle + \langle u, (J + \mu_i I)^{m_i} w'_2 \rangle \\ &= \langle u, w_1 \rangle + \langle (-1)^{m_i} (J - \mu_i I)^{m_i} u, w'_2 \rangle \\ &= \langle u, w_1 \rangle + 0 \\ &= \langle u, w_1 \rangle. \end{aligned}$$

It follows that $\ker(J - \mu_i I)^{m_i}$ and $\ker(J + \mu_i I)^{m_i}$ are complementary null subspaces (of each other).

Similarly, one may show that $\ker(J^2 + 2a_i J + (a_i^2 + b_i^2)I)$ and $\ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)$ are likewise complementary. $\square$

From these two propositions it now follows that the minimal polynomial of J can be written in the form

$$\begin{aligned} p(t) &= \prod_{i=1}^p (t^2 - 2a_i t + (a_i^2 + b_i^2))^{k_i} (t^2 + 2a_i t + (a_i^2 + b_i^2))^{k_i} \\ &\quad \cdot \prod_{i=1}^q (t^2 + \lambda_i^2)^{l_i} \cdot \prod_{i=1}^r (t - \mu_i)^{m_i} (t + \mu_i)^{m_i} \cdot t^s \end{aligned} \tag{2.1}$$

where $a_i \neq 0 \neq b_i$, $\lambda_i > 0$, $\mu_i \neq 0$, and all $a_i, b_i, \lambda_i, \mu_i$ are real.

Proposition 2.3 $\ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)^{k_i} \oplus \ker(J^2 + 2a_i J + (a_i^2 + b_i^2)I)^{k_i}$ for $1 \leq i \leq p$, $\ker(J^2 + \lambda_i^2 I)^{l_i}$ for $1 \leq i \leq q$, $\ker(J - \mu_i I)^{m_i} \oplus \ker(J + \mu_i I)^{m_i}$ for $1 \leq i \leq r$, and $\ker J^s$ are J -invariant, mutually orthogonal, nondegenerate subspaces of V .

Proof: These are rather straightforward calculations. As an example, we show that each $\ker(J^2 + \lambda_i^2 I)^{l_i}$ is nondegenerate.

Suppose a vector $u \in \ker(J^2 + \lambda_i^2 I)^{l_i}$ satisfies $\langle u, v \rangle = 0$ for all $v \in \ker(J^2 + \lambda_i^2 I)^{l_i}$. Let

$$\begin{aligned} W = & \bigoplus_{i=1}^p (\ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)^{k_i} \oplus \ker(J^2 + 2a_i J + (a_i^2 + b_i^2)I)^{k_i}) \\ & \oplus \bigoplus_{i=1}^r (\ker(J - \mu_i I)^{m_i} \oplus \ker(J + \mu_i I)^{m_i}) \\ & \oplus \bigoplus_{j \neq i} \ker(J^2 + \lambda_j^2 I)^{l_j} \oplus \ker J^s. \end{aligned}$$

Then $(J^2 + \lambda_i^2 I)^{l_i} : W \rightarrow W$ is nonsingular, hence surjective. This and $\langle u, (J^2 + \lambda_i^2 I)^{l_i} w \rangle = \langle (J^2 + \lambda_i^2 I)^{l_i} u, w \rangle = 0$ for all $w \in W$ imply that $\langle u, w \rangle = 0$ for all $w \in W$, thus $\langle u, v \rangle = 0$ for all $v \in V$, so $u = 0$ since V is nondegenerate. Consequently $\ker(J^2 + \lambda_i^2 I)^{l_i}$ is also nondegenerate. $\square$

It now follows that J can be decomposed as an orthogonal direct sum of skewadjoint operators on each of these subspaces:

$$\begin{aligned} & \ker(J^2 - 2a_i J + (a_i^2 + b_i^2)I)^{k_i} \oplus \ker(J^2 + 2a_i J + (a_i^2 + b_i^2)I)^{k_i} \\ & \hspace{25em} \text{for } 1 \leq i \leq p; \\ & \ker(J^2 + \lambda_i^2 I)^{l_i} \quad \text{for } 1 \leq i \leq q; \\ & \ker(J - \mu_i I)^{m_i} \oplus \ker(J + \mu_i I)^{m_i} \quad \text{for } 1 \leq i \leq r; \\ & \ker J^s. \end{aligned}$$

Thus it will suffice to classify skewadjoint operators of each type.

3 Minimal polynomial $(t^2 - 2at + (a^2 + b^2))^k (t^2 + 2at + (a^2 + b^2))^k$

If J is such a skewadjoint operator on V , then we can also consider J as a skewadjoint operator on the complexification $U = V^{\mathbb{C}}$ of V . Here the inner

product is extended by bilinearity in the usual way.

$$\langle v_1 + iw_1, v_2 + iw_2 \rangle = \langle v_1, v_2 \rangle - \langle w_1, w_2 \rangle + i(\langle w_1, v_2 \rangle + \langle v_1, w_2 \rangle)$$

The minimal polynomial of J factors over $\mathbb{C}$ as

$$(t - (a + ib))^k (t - (a - ib))^k (t + (a + ib))^k (t + (a - ib))^k.$$

Then there exists a vector $u + iv \in U$ such that $(J - (a + ib)I)^{k-1}(u + iv) \neq 0$ and $(J - (a + ib)I)^k(u + iv) = 0$. Since $\operatorname{Re}(J - (a + ib)I)^{k-1}(u + iv)$ and $\operatorname{Im}(J - (a + ib)I)^{k-1}(u + iv)$ lie in $\ker(J^2 - 2aJ + (a^2 + b^2)I)^k$, there exist $w, x \in \ker(J^2 + 2aJ + (a^2 + b^2)I)^k$ such that

$$\langle (J - (a + ib)I)^{k-1}(u + iv), w + ix \rangle = 2 \quad (3.1)$$

by Proposition 2.2. Now we need the following result.

Lemma 3.1 *If $u + iv \in \ker(J - (a + ib)I)^k$ and $w + ix \in \ker(J + (a - ib)I)^k$, then $\langle u + iv, w + ix \rangle = 0$.*

Proof: Since

$$(J + (a + ib)I)^k : \ker(J + (a - ib)I)^k \rightarrow \ker(J + (a - ib)I)^k$$

is nonsingular, there exists $w' + ix' \in \ker(J + (a - ib)I)^k$ such that

$$(J + (a + ib)I)^k(w' + ix') = w + ix.$$

Thus,

$$\begin{aligned} \langle u + iv, w + ix \rangle &= \langle u + iv, (J + (a + ib)I)^k(w' + ix') \rangle \\ &= \langle (-1)^k (J - (a + ib)I)^k(u + iv), w' + ix' \rangle \\ &= \langle 0, w' + ix' \rangle \\ &= 0. \end{aligned}$$

□

Write $w + ix = w_1 + ix_1 + w_2 + ix_2$ with $w_1 + ix_1 \in \ker(J + (a + ib)I)^k$ and $w_2 + ix_2 \in \ker(J + (a - ib)I)^k$. Then from (3.1) and Lemma 3.1 we obtain

$$\begin{aligned} 2 &= \langle (J - (a + ib)I)^{k-1}(u + iv), w + ix \rangle \\ &= \langle (-(a + ib)I)^{k-1}(u + iv), w_1 + ix_1 + w_2 + ix_2 \rangle \\ &= \langle (J - (a + ib)I)^{k-1}(u + iv), w_1 + ix_1 \rangle \end{aligned}$$

This implies that we may assume that

$$w + ix \in \ker (J + (a + ib)I)^k$$

in (3.1). Since $w - ix \in \ker (J + (a - ib)I)^k$, we have *via* Lemma 3.1

$$\langle (J - (a + ib)I)^h (u + iv), (J + (a - ib)I)^j (w - ix) \rangle = 0 \quad (3.2)$$

for all nonnegative integers h, j .

Also, we can find complex numbers $a_1, a_2, \dots, a_{k-1}$ such that

$$\begin{aligned} & \langle (u + iv) + a_1(J - (a + ib)I)(u + iv) + \dots \\ & \quad + a_{k-1}(J - (a + ib)I)^{k-1}(u + iv), w + ix \rangle = 0, \\ & \langle (J - (a + ib)I)(u + iv) + a_1(J - (a + ib)I)^2(u + iv) + \dots \\ & \quad + a_{k-2}(J - (a + ib)I)^{k-1}(u + iv), w + ix \rangle = 0, \\ & \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \\ & \langle (J - (a + ib)I)^{k-2}(u + iv) + a_1(J - (a + ib)I)^{k-1}(u + iv), w + ix \rangle = 0. \end{aligned}$$

Replacing $u + iv$ by

$$(u + iv) + a_1(J - (a + ib)I)(u + iv) + \dots + a_{k-1}(J - (a + ib)I)^{k-1}(u + iv)$$

we have

$$\langle (J - (a + ib)I)^h (u + iv), w + ix \rangle = 0 \text{ for } 0 \leq h \leq k - 2. \quad (3.3)$$

Note that (3.1) and (3.2) continue to hold as well.

Let $u_1 = u$ and $v_1 = v$ and set

$$(J - (a + ib)I)^h (u_1 + iv_1) = u_{h+1} + iv_{h+1} \text{ for } 1 \leq h \leq k - 1. \quad (3.4)$$

Then

$$\begin{aligned} Ju_h &= au_h - bv_h + u_{h+1} \\ Jv_h &= bu_h + av_h + v_{h+1} \end{aligned} \quad \text{for } 1 \leq h \leq k - 1 \quad (3.5)$$

$$\begin{aligned} Ju_k &= au_k - bv_k \\ Jv_k &= bu_k + av_k \end{aligned} \quad (3.6)$$

Now (3.1) implies that

$$\begin{aligned} (J + (a + ib)I)^{k-1} (w + ix) &\neq 0, \\ (J + (a + ib)I)^k (w + ix) &= 0. \end{aligned}$$

Let $w_1 = w$ and $x_1 = x$ and set

$$(J + (a + ib)I)^h(w_1 + ix_1) = w_{h+1} + ix_{h+1} \text{ for } 1 \leq h \leq k-1. \quad (3.7)$$

Then

$$\begin{aligned} Jw_h &= -aw_h + bx_h + w_{h+1} \\ Jx_h &= -bw_h - ax_h + x_{h+1} \end{aligned} \quad \text{for } 1 \leq h \leq k-1 \quad (3.8)$$

$$\begin{aligned} Jw_k &= -aw_k + bx_k \\ Jx_k &= -bw_k - ax_k \end{aligned} \quad (3.9)$$

By (3.1), (3.2), and (3.3),

$$\begin{aligned} &\langle (J - (a + ib)I)^h(u_1 + iv_1), (-1)^j(J + (a + ib)I)^j(w_1 + ix_1) \rangle \\ &= \begin{cases} 2 & \text{if } h + j = k - 1, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

$$\langle (J - (a + ib)I)^h(u_1 + iv_1), (-1)^j(J + (a - ib)I)^j(w_1 - ix_1) \rangle = 0$$

for all nonnegative integers h, j . This implies that

$$\begin{aligned} \langle u_{h+1} + iv_{j+1}, (-1)^j(w_{j+1} + ix_{j+1}) \rangle &= \begin{cases} 2 & \text{if } h + j = k - 1, \\ 0 & \text{otherwise,} \end{cases} \\ \langle u_{h+1} + iv_{h+1}, w_{j+1} - ix_{j+1} \rangle &= 0 \end{aligned}$$

for all nonnegative integers h, j . Thus we have all inner products zero except

$$\begin{aligned} \langle u_h, w_j \rangle &= (-1)^{j-1} \\ \langle v_h, x_j \rangle &= (-1)^j \end{aligned} \quad \text{if } h + j = k + 1.$$

(Recall that $\ker(J^2 - 2aJ + (a^2 + b^2)I)$ and $\ker(J^2 + 2aJ + (a^2 + b^2)I)$ are null subspaces.) Note that $\{u_1, v_1, \dots, u_k, v_k, w_1, x_1, \dots, w_k, x_k\}$ spans a J -invariant, nondegenerate subspace. Hence there exists a J -invariant, nondegenerate subspace N of V such that

$$\llbracket u_1, v_1, \dots, u_k, v_k, w_1, x_1, \dots, w_k, x_k \rrbracket \oplus N$$

is an orthogonal direct sum. On the subspace $\llbracket u_1, v_1, \dots, w_k, x_k \rrbracket$, the matrix of J is $2k \times 2k$ of the following form.

$$\begin{bmatrix} a & b & & & & & & & & \\ -b & a & & & & & & & & \\ 1 & 0 & a & b & & & & & & \\ 0 & 1 & -b & a & & & & & & \\ & & & \ddots & & & & & & \\ & & & & a & b & & & & \\ & & & & -b & a & & & & \\ & & & & 1 & 0 & a & b & & \\ & & & & 0 & 1 & -b & a & & \\ & & & & & & -a & -b & & \\ & & & & & & b & -a & & \\ & & & & & & 1 & 0 & -a & -b \\ & & & & & & 0 & 1 & b & -a \\ & & & & & & & & \ddots & \\ & & & & & & & & & -a & -b \\ & & & & & & & & & b & -a \\ & & & & & & & & & 1 & 0 & -a & -b \\ & & & & & & & & & 0 & 1 & b & -a \end{bmatrix}$$

Continuing this process, we get an orthogonal direct sum of J -invariant subspaces V_h such that on each of them, J has a matrix of the preceding form but of size $2r \times 2r$ for some r with $1 \leq r \leq k$, with respect to a basis of the specified type.

4 Minimal polynomial $(t^2 + \lambda^2)^l$

Let J be a skewadjoint operator on V with such a minimal polynomial.

Lemma 4.1 *There exists a vector $v \in V$ such that $v, (J^2 + \lambda^2 I)v, \dots, (J^2 + \lambda^2 I)^{l-1}v$ are linearly independent and $\langle v, (J^2 + \lambda^2 I)^{l-1}v \rangle \neq 0$.*

Proof: Since the minimal polynomial of J is $(t^2 + \lambda^2)^l$, there exists $v \in V$ such that $v, (J^2 + \lambda^2 I)v, \dots, (J^2 + \lambda^2 I)^{l-1}v$ are linearly independent.

If $\langle v, (J^2 + \lambda^2 I)^{l-1}v \rangle = 0$, then there exists $w \in V$ such that $\langle w, (J^2 + \lambda^2 I)^{l-1}v \rangle \neq 0$. If $\langle w, (J^2 + \lambda^2 I)^{l-1}w \rangle \neq 0$, then we simply replace v with w and verify that we have achieved the desired result; otherwise, replace v with $v + w$ and verify that this works. $\square$

So we may start with $v \in V$ as in the Lemma. Set $(J + i\lambda I)^l v = u_1 + iv_1$ and observe that $(u_1 + iv_1), (J - i\lambda I)(u_1 + iv_1), \dots, (J - i\lambda I)^{l-1}(u_1 + iv_1)$ are linearly independent over $\mathbb{C}$ and that $(J - i\lambda I)^l(u_1 + iv_1) = (J^2 + \lambda^2 I)^l v = 0$. Let

$$(J - i\lambda I)^h (u_1 + iv_1) = u_{h+1} + iv_{h+1} \quad \text{for } 0 \leq h \leq l-1.$$

First off, we have

$$\langle u_h + iv_h, u_j + iv_j \rangle = 0 \quad \text{for } 1 \leq h, j \leq l$$

since

$$\begin{aligned} \langle u_h + iv_h, u_j + iv_j \rangle &= \langle (J - i\lambda I)^{h-1}(J + i\lambda I)^l v, (J - i\lambda I)^{j-1}(J + i\lambda I)^l v \rangle \\ &= (-1)^l \langle (J - i\lambda I)^{h-1}(J - i\lambda I)^l (J + i\lambda I)^l v, (J - i\lambda I)^{j-1} v \rangle \\ &= (-1)^l \langle (J - i\lambda I)^{h-1}(J^2 + \lambda^2 I)^l v, (J - i\lambda I)^{j-1} v \rangle \\ &= 0. \end{aligned}$$

This implies that

$$\begin{aligned} \langle u_h, u_j \rangle - \langle v_h, v_j \rangle &= 0 \\ \langle u_h, v_j \rangle + \langle v_h, u_j \rangle &= 0 \end{aligned} \quad \text{for } 1 \leq h, j \leq l. \quad (4.1)$$

Next, we have

$$\begin{aligned} \langle u_l + iv_l, u_h - iv_h \rangle &= \langle (J - i\lambda I)^{l-1}(J + i\lambda I)^l v, (J + i\lambda I)^{h-1}(J - i\lambda I)^l v \rangle \\ &= (-1)^{h-1} \langle (J - i\lambda I)^{l+h-2}(J + i\lambda I)^l v, (J - i\lambda I)^l v \rangle \\ &= (-1)^{h-1} \langle (J - i\lambda I)^{h-2}(J^2 + \lambda^2 I)^l v, (J - i\lambda I)^l v \rangle \end{aligned}$$

whence

$$\langle u_l + iv_l, u_h - iv_h \rangle = 0 \quad \text{for } 2 \leq h \leq l. \quad (4.2)$$

Note that $J^2(J^2 + i\lambda I)^{l-1}v = -\lambda^2(J^2 + i\lambda^2 I)^{l-1}v$. We use this in the following computation.

Lemma 4.2 *If l is odd, then $\langle u_1 + iv_1, u_l - iv_l \rangle$ is a nonzero real number, and if l is even then it is nonzero pure imaginary. Moreover, $\langle u_h + iv_h, u_j - iv_j \rangle$ is real when $h + j$ is even and pure imaginary when $h + j$ is odd.*

Proof: We compute

$$\begin{aligned}
& \langle u_1 + iv_1, u_l - iv_l \rangle \\
&= \langle (J + i\lambda I)^l v, (J + i\lambda I)^{l-1} (J - i\lambda I)^l v \rangle \\
&= -\langle (J + i\lambda I)^{l+1} v, (J^2 + \lambda^2 I)^{l-1} v \rangle \\
&= -\langle \left(J^{l+1} + \binom{l+1}{1} (i\lambda) J^l + \dots + \binom{l+1}{l} (i\lambda)^l J + (i\lambda)^{l+1} I \right) v, (J^2 + \lambda^2 I)^{l-1} v \rangle.
\end{aligned}$$

If l is odd,

$$\begin{aligned}
\langle u_1 + iv_1, u_l - iv_l \rangle &= -\langle \left(J^{l+1} - \binom{l+1}{2} \lambda^2 J^{l-1} + \binom{l+1}{4} \lambda^4 J^{l-3} - \dots \right. \\
&\quad \left. + (-1)^{\frac{l+1}{2}} \lambda^{l+1} I \right) v, (J^2 + \lambda^2 I)^{l-1} v \rangle \\
&= -\langle \left((-1)^{\frac{l+1}{2}} \lambda^{l+1} - \binom{l+1}{2} \lambda^2 (-1)^{\frac{l-1}{2}} \lambda^{l-1} + \dots \right. \\
&\quad \left. + (-1)^{\frac{l+1}{2}} \lambda^{l+1} \right) v, (J^2 + \lambda^2 I)^{l-1} v \rangle \\
&= (-1)^{\frac{l+3}{2}} \lambda^{l+1} \left(1 + \binom{l+1}{2} + \binom{l+1}{4} + \dots \right. \\
&\quad \left. + \binom{l+1}{l-1} + 1 \right) \langle v, (J^2 + \lambda^2 I)^{l-1} v \rangle
\end{aligned}$$

which is a nonzero real number. If l is even, an analogous computation yields a nonzero pure imaginary number. The rest is done similarly. $\square$

Now assume that l is odd and consider a vector of the form

$$w + ix = (u_1 + iv_1) + \alpha_1(u_2 + iv_2) + \dots + \alpha_{l-1}(u_l + iv_l).$$

Then we can determine the (complex) coefficients α_i such that

$$\langle w + ix, (J + i\lambda I)^h (w - ix) \rangle = 0 \quad \text{for } 0 \leq h \leq l-2.$$

Indeed, consider

$$\begin{aligned}
& \langle w + ix, (J + i\lambda I)^{l-2} (w - ix) \rangle \\
&= \langle (u_1 + iv_1) + \alpha_1(u_2 + iv_2) + \dots + \alpha_{l-1}(u_l + iv_l), \\
&\quad (u_{l-1} - iv_{l-1}) + \bar{\alpha}_1(u_l - iv_l) \rangle \\
&= \langle u_1 + iv_1, u_{l-1} - iv_{l-1} \rangle + \alpha_1 \langle u_2 + iv_2, u_{l-1} - iv_{l-1} \rangle \\
&\quad + \bar{\alpha}_1 \langle u_1 + iv_1, u_l - iv_l \rangle.
\end{aligned}$$

Since $\langle u_1 + iv_1, u_{l-1} - iv_{l-1} \rangle$ is pure imaginary and $\langle u_2 + iv_2, u_{l-1} - iv_{l-1} \rangle = -\langle u_1 + iv_1, u_l - iv_l \rangle$ is real, we can find a pure imaginary α_1 such that

$$\langle w + ix, (J + i\lambda I)^{l-2} (w - ix) \rangle = 0.$$

At the next step, there are six such inner product terms (with coefficients) which are all real. We obtain a real α_2 such that

$$\langle w + ix, (J + i\lambda I)^{l-3}(w - ix) \rangle = 0.$$

Continuing inductively, we find all the desired coefficients.

Replacing the previous $u_1 + iv_1$ with $w + ix$ so determined, using (4.1), (4.2), and Lemma 4.2, and rescaling, we have

$$\begin{aligned} \langle u_h + iv_h, u_j + iv_j \rangle &= 0 \quad \text{for } 1 \leq h, j \leq l; \\ \langle u_h + iv_h, u_j - iv_j \rangle &= \begin{cases} (-1)^j(\pm 2) & \text{if } h + j = l + 1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

In other words, all inner products among the u_h and v_j are zero except

$$\langle u_h, u_j \rangle = \langle v_h, v_j \rangle = (-1)^j(\pm 1) \quad \text{if } h + j = l + 1.$$

Since $(J - i\lambda I)(u_h + iv_h) = u_{h+1} + iv_{h+1}$ for $1 \leq h \leq l-1$ and $(J - i\lambda I)(u_l + iv_l) = 0$, the matrix of J on $\llbracket u_1, v_1, \dots, u_l, v_l \rrbracket$ is $2l \times 2l$ of this form.

$$\begin{bmatrix} 0 & \lambda & & & & \\ -\lambda & 0 & & & & \\ 1 & 0 & 0 & \lambda & & \\ 0 & 1 & -\lambda & 0 & & \\ & & & \ddots & & \\ & & & & 0 & \lambda \\ & & & & -\lambda & 0 \\ & & & & 1 & 0 & 0 & \lambda \\ & & & & 0 & 1 & -\lambda & 0 \end{bmatrix}$$

If l is even, the condition on inner products of the basis vectors becomes

$$\langle v_h, u_j \rangle = -\langle u_h, v_j \rangle = (-1)^j(\pm 1) \quad \text{if } h + j = l + 1$$

with all others vanishing as before, while the $2l \times 2l$ form of the matrix of J remains the same.

We finish by continuing this process as at the end of the preceding section, with $1 \leq r \leq l$ now.

5 Minimal polynomial $(t - \mu)^m(t + \mu)^m$

Let $J : V \rightarrow V$ be a skewadjoint operator with such a minimal polynomial. Then $V = \ker(J - \mu I)^m \oplus \ker(J + \mu I)^m = W_1 \oplus W_2$ where W_1 and W_2 are complementary null subspaces by Propositions 2.1 and 2.2. There exists $v \in W_1$ such that $v, (J - \mu I)v, \dots, (J - \mu I)^{m-1}v$ are linearly independent and $(J - \mu I)^m v = 0$. Then we can choose $w \in W_2$ such that $\langle (J - \mu I)^{m-1}v, w \rangle = 1$ and there exist real numbers $a_1, \dots, a_{m-1}$ such that

$$\begin{aligned} \langle v + a_1(J - \mu I)v + \dots + a_{m-1}(J - \mu I)^{m-1}v, w \rangle &= 0 \\ \langle (J - \mu I)v + a_1(J - \mu I)^2v + \dots + a_{m-2}(J - \mu I)^{m-1}v, w \rangle &= 0 \\ \vdots & \qquad \qquad \qquad \vdots \qquad \qquad \qquad \vdots \\ \langle (J - \mu I)^{m-2}v + a_1(J - \mu I)^{m-1}v, w \rangle &= 0 \end{aligned}$$

If we replace v by $v + a_1(J - \mu I)v + \dots + a_{m-1}(J - \mu I)^{m-1}v$ and set $v_1 = v$, $v_2 = (J - \mu I)v_1$, $\dots$, $v_m = (J - \mu I)v_{m-1}$, and $w_1 = w$, $w_2 = (J + \mu I)w_1$, $\dots$, $w_m = (J + \mu I)w_{m-1}$, then all inner products are zero except

$$\begin{aligned} \langle v_i, w_j \rangle &= \langle (J - \mu I)^{i-1}v, (J + \mu I)^{j-1}w \rangle \\ &= \langle (-1)^{j-1}(J - \mu I)^{i+j-2}v, w \rangle \\ &= (-1)^{j-1} \quad \text{if } i + j = m + 1. \end{aligned}$$

We also have a J -invariant, nondegenerate subspace N such that $V = \llbracket v_1, \dots, v_m, w_1, \dots, w_m \rrbracket \oplus N$ is an orthogonal direct sum. With respect to the obvious basis, the matrix of J on the first subspace is block-diagonal

$$\begin{bmatrix} \mu & & & & & & \\ 1 & \mu & & & & & \\ & 1 & \ddots & & & & \\ & & 1 & \mu & & & \\ & & & -\mu & & & \\ & & & 1 & -\mu & & \\ & & & & 1 & \ddots & \\ & & & & & 1 & -\mu \end{bmatrix}$$

with two elementary Jordan diagonal blocks, each $m \times m$.

We finish as before, with $1 \leq r \leq m$ now.

6 Minimal polynomial t^s

Let J be such a skewadjoint operator. Then there exists a subspace V_1 of V such that $V = \ker J^{s-1} \oplus V_1$, whence $J^{s-1}V = J^{s-1}V_1$. Since $J^{s-1}V_1$ is a null subspace, there exists a complementary null subspace W_1 and a subspace N which is orthogonal to their direct sum so that $V = (J^{s-1}V_1 \oplus W_1) \oplus N$. Then $J^{s-1}V = J^{s-1}V_1 = J^{s-1}W_1 \oplus J^{s-1}N$. Since W_1 is a complementary null subspace to $J^{s-1}V_1$, this implies that $J^{s-1}V_1 = J^{s-1}W_1$ and that $J^{s-1}N \subseteq J^{s-1}W_1$. Thus we may assume that $W_1 = V_1$ and we have

$$V = J^{s-1}V_1 \oplus V_1 \oplus N$$

where $J^{s-1}V_1$ and V_1 are complementary null subspaces (of each other) and N is orthogonal to their (direct) sum.

Firstly, assume that s is even. Consider $J^{s-1}v_1$ for a nonzero vector in V_1 . Then there exists $w_1 \in V_1$ such that $\langle J^{s-1}v_1, w_1 \rangle = 1$ and v_1 and w_1 are linearly independent.

Lemma 6.1 *$\llbracket v_1, \dots, J^{s-1}v_1, w_1, \dots, J^{s-1}w_1 \rrbracket$ is a J -invariant, nondegenerate, $2s$ -dimensional subspace of V .*

Proof: Suppose $a_1v_1 + \dots + a_sJ^{s-1}v_1 + b_1w_1 + \dots + b_sJ^{s-1}w_1 = 0$ for real a_i, b_i . Applying J^{s-1} to this equation, we obtain $a_1J^{s-1}v_1 + b_1J^{s-1}w_1 = 0$. Then $\langle a_1J^{s-1}v_1 + b_1J^{s-1}w_1, w_1 \rangle = a_1 = 0$ whence $b_1 = 0$. Continuing this process, all the coefficients vanish so our list is a basis and the subspace is $2s$ -dimensional.

If $\langle a_1v_1 + \dots + J^{s-1}v_1 + b_1w_1 + \dots + J^{s-1}w_1, u \rangle = 0$ for all u in our subspace with real a_i, b_i , then replacing u by $J^{s-1}w_1$ yields $a_1 = 0$. Continuing in this way, all the coefficients vanish again; this time it shows that our subspace is nondegenerate. $\square$

Next, we can find real numbers a_i, b_j such that if $v = v_1 + a_1Jv_1 + a_2J^2v_1 + a_3J^3v_1 + \dots + a_{s-1}J^{s-1}v_1 + b_1Jw_1 + b_3J^3w_1 + b_5J^5w_1 + \dots + b_{s-1}J^{s-1}w_1$, then

$$\langle v, J^h v \rangle = \langle v, J^h w_1 \rangle = 0 \quad \text{for } 0 \leq h \leq s-2.$$

Replacing v_1 by this v , we may assume that

$$\begin{aligned} \langle v_1, J^h v_1 \rangle &= 0 & \text{for } 0 \leq h, \\ \langle v_1, J^h w_1 \rangle &= 0 & \text{for } 0 \leq h \leq s-2, \\ \langle J^{s-1}v_1, w_1 \rangle &= 1. \end{aligned}$$

Now we can find real c_i such that if $w = w_1 + c_1 J v_1 + c_3 J^3 v_1 + c_5 J^5 v_1 + \cdots + c_{s-1} J^{s-1} v_1$, then $\langle w, J^h w \rangle = 0$ for all $h \geq 0$. Replacing w_1 by this w , we may assume that all inner products $\langle v_1, J^h v_1 \rangle$, $\langle v_1, J^h w_1 \rangle$, $\langle w_1, J^h w_1 \rangle$ vanish except $\langle J^{s-1} v_1, w_1 \rangle = 1$. Setting $v_h = J^{h-1} v_1$ and $w_h = J^{h-1} w_1$, the matrix of J on the basis $\{v_1, \dots, v_s, w_1, \dots, w_s\}$ is $2s \times 2s$ of the form

$$\begin{bmatrix} 0 & & & & & & & & & & \\ 1 & 0 & & & & & & & & & \\ & 1 & \ddots & & & & & & & & \\ & & & 1 & 0 & & & & & & \\ & & & & 1 & 0 & & & & & \\ & & & & & 0 & & & & & \\ & & & & & 1 & 0 & & & & \\ & & & & & & 1 & \ddots & & & \\ & & & & & & & 1 & 0 & & \\ & & & & & & & & 1 & 0 & \\ & & & & & & & & & 1 & 0 \end{bmatrix}$$

and all inner products vanish except $\langle v_i, w_j \rangle = (-1)^{j-1}$ if $i + j = s + 1$.

Lastly, assume that s is odd. Consider $J^{s-1} v_1$ for $v_1 \in V$. Then there exists $w_1 \in V$ such that $\langle J^{s-1} v_1, w_1 \rangle = \frac{1}{2}$ and $\langle J^{s-1}(v_1 + w_1), v_1 + w_1 \rangle = 1$. We proceed by replacing v_1 by $v_1 + w_1$. As before, we then have that $\llbracket v_1, J v_1, \dots, J^{s-1} v_1 \rrbracket$ is a J -invariant, nondegenerate, s -dimensional subspace of V . Also, we can find real numbers $a_2, a_4, \dots, a_{s-1}$ such that

$$\langle v_1 + a_2 J^2 v_1 + \cdots + a_{s-1} J^{s-1} v_1, J^h (v_1 + a_2 J^2 v_1 + \cdots + a_{s-1} J^{s-1} v_1) \rangle = 0$$

for $0 \leq h \leq s-2$. Replacing the current v_1 by $v_1 + a_2 J^2 v_1 + \cdots + a_{s-1} J^{s-1} v_1$, we obtain

$$\begin{aligned} \langle v_1, J^h v_1 \rangle &= 0 \quad \text{for } 0 \leq h \leq s-2, \\ \langle v_1, J^{s-1} v_1 \rangle &= 1. \end{aligned}$$

Set $v_h = J^{h-1} v_1$ for $1 \leq h \leq s$. The the matrix of J with respect to the basis $\{v_1, v_2, \dots, v_s\}$ is

$$\begin{bmatrix} 0 & & & & & \\ 1 & 0 & & & & \\ & 1 & \ddots & & & \\ & & & 1 & 0 & \\ & & & & 1 & 0 \end{bmatrix}$$

with all inner products of basis vectors vanishing except $\langle v_i, v_j \rangle = (-1)^{j-1}$ if $i + j = s + 1$.

We finish almost as usual, with $1 \leq r \leq s$, but $r \times r$ blocks now. This completes the classification.

References

- [1] G. Birkhoff and S. Mac Lane, *A Survey of Modern Algebra*, 3rd ed. New York: Macmillan, 1965.
- [2] L. A. Cordero and P. E. Parker, Pseudoriemannian 2-step nilpotent Lie groups, Santiago-Wichita preprint DGS/CP4, 1999. arXiv: math.DG/9905188
- [3] P. Eberlein, Geometry of 2-step nilpotent groups with a left invariant [*sic*] metric, *Ann. scient. Éc. Norm. Sup.* **27** (1994) 611–660.
- [4] R. Horn and C. Johnson, *Matrix Analysis*. Cambridge: U. P., 1985.
- [5] C. Jang and K. Park, Conjugate points on 2-step nilpotent groups, *Geom. Dedicata* **79** (2000) 65–80.
- [6] C. Jang and P. E. Parker, Conjugate loci of pseudoriemannian 2-step nilpotent Lie groups with nondegenerate center, Wichita-Ulsan preprint DGS/JP1, 2003.
- [7] B. O’Neill, *Semi-Riemannian Geometry*. New York: Academic Press, 1983.